

Medical Science

To Cite:

Kościan J, Mitkowska M, Dryżałowska Z, Obrębski M, Szczerkowska K, Chwiejczak J, Górny A, Seroka A, Rybicka M, Młynarska J, Wójcik A. Binge Eating Disorder and Loss-of-control Eating following bariatric surgery: An update on research. *Medical Science* 2024; 28: e72ms3375 doi: <https://doi.org/10.54905/disssi.v28i149.e72ms3375>

Authors' Affiliation:

¹Independent Public Hospital them. prof. W. Orłowski Medical Centre of Postgraduate Education, Czerniakowska 231, 00-401, Warsaw, Poland

²HCP Medical Centre, Poznan, Poland

³University Clinical Centre of the Medical University of Warsaw, Żwirki i Wigury 61, 02-091 Warsaw, Poland

⁴Military Institute of Medicine - National Research Institute, Warsaw, Poland

⁵Memorial Bielański Hospital, Warsaw, Poland

⁶Prague Hospital of Lord's Transfiguration, Warsaw, Poland

⁷The National Institute of Medicine of the Ministry of Interior and Administration, Warsaw, Poland

⁸Saint Anna Hospital of Trauma Surgery, Warsaw, Poland

⁹Warsaw Southern Hospital, Rotmistrza Witolda Pileckiego 99, 02-781 Warszawa, Poland

*Corresponding Author

Independent Public Hospital them. prof. W. Orłowski Medical Centre of Postgraduate Education, Czerniakowska 231, 00-401, Warsaw, Poland
Email: jankoscian@gmail.com

Contact List

Jan Kościan	jankoscian@gmail.com
Maria Mitkowska	mmitkowska1@gmail.com
Zuzanna Dryżałowska	zdryzalowska@gmail.com
Michał Obrębski	ninurt@yahoo.com
Karolina Szczerkowska	kszczerkowska@wim.mil.pl
Justyna Chwiejczak	justynachwiejczak@gmail.com
Aleksander Górny	alexo001444@gmail.com
Anna Seroka	ania.seroka1@gmail.com
Maria Rybicka	rybickamaria97@gmail.com
Julita Młynarska	julitamartamlynarska@gmail.com
Anna Wójcik	awannawojcikaw@gmail.com

ORCID List

Jan Kościan	0009-0005-2769-8714
Maria Mitkowska	0009-0003-9243-2045
Zuzanna Dryżałowska	0009-0005-5182-9147
Michał Obrębski	0009-0000-0420-5853
Karolina Szczerkowska	0009-0003-5995-0633
Justyna Chwiejczak	0009-0005-3269-6289
Aleksander Górny	0009-0009-4301-7842
Anna Seroka	0009-0001-4359-6698
Maria Rybicka	0009-0005-5965-5938
Julita Młynarska	0009-0002-4268-065X
Anna Wójcik	0009-0003-2208-046X

Peer-Review History

Received: 15 April 2024

Reviewed & Revised: 19/April/2024 to 26/June/2024

Accepted: 03 July 2024

Published: 07 July 2024

Peer-review Method

External peer-review was done through double-blind method.

Medical Science

pISSN 2321-7359; eISSN 2321-7367



© The Author(s) 2024. Open Access. This article is licensed under a [Creative Commons Attribution License 4.0 \(CC BY 4.0\)](https://creativecommons.org/licenses/by/4.0/), which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>.



Binge Eating Disorder and Loss-of-control Eating following bariatric surgery: An update on research

Jan Kościan^{1*}, Maria Mitkowska², Zuzanna Dryżałowska¹, Michał Obrębski³, Karolina Szczerkowska⁴, Justyna Chwiejczak⁵, Aleksander Górny⁶, Anna Seroka⁷, Maria Rybicka³, Julita Młynarska⁸, Anna Wójcik⁹

ABSTRACT

Bariatric surgery remains a leading treatment for obesity due to its effectiveness in reducing food intake. However, outcomes may be influenced by various factors, and some patients fail to achieve desired weight reduction. Binge eating disorders or loss of control eating are the subject of interest as there is growing evidence indicating their impact on surgery outcomes. This paper aims to compare recent findings with the existing reviews on eating pathologies in patients undergoing bariatric surgery and summarize the newest discoveries addressing this topic. As patients continue to experience suboptimal weight loss or significant weight regain, there is a growing need to understand the underlying mechanisms of this process better. The comparison of previous reviews and recent findings underscores that greater emphasis should be placed on long-term observation alongside standard pre-surgery preparation. With new medical phenomena emerging, such as nibbling, grazing, and picking, clarification and unification of diagnostic processes should be the aim of future research. All researchers agree that self-monitoring plays a vital role in the post-surgery period. It should be considered a gold standard in all patients undergoing bariatric surgery, enabling the prompt addressing of the needs as they arise.

Keywords: Bariatric surgery, binge eating disorder, loss-of-control eating

1. INTRODUCTION

Overweight, defined as BMI > 25kg/m², and obesity, defined as BMI > 30kg/m², have been proven to be associated with a higher incidence of various conditions such as hypertension, cardiovascular diseases, diabetes mellitus, renal and liver

failure, mental disorders, and countless other complications. Obesity has been associated with numerous prevalent cancers, such as breast, colorectal, esophageal, kidney, gallbladder, uterine, pancreatic, and liver cancer. The number of obese people has tripled since 1975, and this concern is no longer affecting only developed countries but all nations worldwide. Overweight and obesity are diagnosed among around 40% of the global population. According to the World Obesity Federation, this number is expected to grow to over 50% by 2035.

Currently, obesity alone affects approximately 15% of the population and is estimated to reach even 24% by 2035, impacting nearly 2 billion people, including adults, adolescents, and children. It is worth noting that the most rapid increase in obesity cases is expected among boys and girls – between 2020 and 2035, the percentage of obese children is expected to double, rising from 10% to 20% for boys and from 8% to 18% for girls (Lobstein et al., 2023). Bariatric surgery continues to stand out as the most effective obesity treatment (Courcoulas et al., 2018). According to data from the International Federation for the Surgery of Obesity and Metabolic Disorders, in 2023, approximately 600,000 such surgeries were performed worldwide (Brown et al., 2023). Weight loss, if maintained, can result mitigate adverse obesity-related comorbidities, thus contributing to prolonging life (Hassapidou et al., 2023; Obeso-Fernández et al., 2023).

However, beyond the surgery itself, many other factors can influence the outcome for patients (Noria et al., 2023; O'Brien et al., 2019; Athanasiadis et al., 2021). One of the obstacles patients have to battle are eating disorders, affecting both pre- and post-surgery period (Law et al., 2023). The most common ones are thought to be maladaptive or dysregulated eating (Kloock et al., 2023). Especially the former and its subtypes – binge eating disorder (BED) and loss-of-control eating (LOCE) – are responsible for poor weight loss results. Poor adherence to post-surgery medical recommendations is the reason why most bariatric programs oblige their participants to undergo a mental health evaluation process before the surgery (Sarwer and Heinberg, 2020). Unfortunately, little to no effort is taken to prevent those dysfunctional eating patterns from manifesting in the months following the procedure.

It is particularly disconcerting, considering that there is a positive correlation between postoperative eating disinhibition and weight regain (Athanasiadis et al., 2021). Binge eating disorder is widely understood that BED is the most prevalent eating disorder occurring prior to surgery, with an incidence of 4-49% (Niego et al., 2007; Opozda et al., 2016). Loss-of-control eating, as defined in DSM-V, is the inability to control what or how much one eats, regardless of the amount of food consumed. Since, after bariatric surgery, ingestion of objectively large amounts of food is limited by new gastric capacity, LOCE is sometimes used in place of BED to better depict the pathological behavior. Evidence shows that up to 39% of patients develop LOCE within two years post-surgery (White et al., 2010).

2. METHOD

This review aimed to consolidate information obtained by previous reviews Opozda et al., (2016), Williams-Kerver et al., (2019), Taba et al., (2021) with a comparison to new articles covering the topic of bariatric surgery and eating disorders. Studies were eligible for inclusion, if they reported mainly on LOCE and BED, both before and after the surgery, and their relation to post-surgery weight outcomes. Studies aiming for systematizing post-surgery recommendations and diagnostic criteria were included. Only studies including adult participants were included. The MEDLINE database was searched via the Pubmed interface up to 30 January 2024. The keywords were (eating) AND (disorder) AND (bariatric) into the database screening only for articles published in the last five years.

3. RESULTS

Out of 382 results, 166 were excluded due to a lack of full articles available. The remaining 216 were screened by the title. Of this, 27 articles covering the topic were evaluated by reading the abstract, then only six studies Allison et al., (2023), Furtado et al., (2023), Lydecker et al., (2019), Ivezaj et al., (2021), Sarwer et al., (2021), Conceição and Goldschmid, (2019) were included, apart from previously mentioned reviews. Additionally, one referenced review was included (Meany et al., 2014). The study selection is shown in (Figure 1). The post-surgical follow-up period ranged from 6 months to 7 years, with final assessment after 12 months in most cases. The results of each latest study are presented in (Table 1).




Figure 1 Study selection flow diagram

Table 1 Recent findings in post-bariatric eating disorders

Authors, year	Sample Size	BED	LOCE	Other findings	Methods
Conceição et al., (2015)	-	-	-	Researchers recommend standardizing diagnostic schemes and assessment tools	-
Furtado et al., (2023)	217	Preoperative BED was present in 19 patients; within two years after the surgery there were 54 individuals screened positive for BED	-	Individuals with BED were 2 times more likely to regain >15% of weight after the surgery	BED, HADS

Ivezaj et al., (2021)	-	-	-	Researchers aimed to compare questionnaires used to assess eating disorders	EDE-BSV, BES, EDDS, EDI, NEQ
-	268 at the start, 186 completed full follow-up	28/268 with history of BED, including 18 who met BED criteria currently. After 12 months, there were no new cases, and only 6/28 met the criteria for BED	-	Patients with subjective binge eating episodes experienced more significant weight loss by 4,1%	SCID-5-RV, EDE-BSV
Lydecker et al., (2021)	168, only with reported LOCE	71 met bariatric BED criteria, and 97 met subthreshold bariatric BED criteria	All 168 reported regular LOCE in 6 months after the surgery	The study also aimed to examine secretive eating – 36,9% (62/168) reported this disorder	EDE-BSV, BDI-II
Sarwer et al., (2021)	300	27 met diagnostic criteria	It was not the focus of the study; however, many reported a feeling of LOCE	Study provides evidence for a coincidence of obesity with other psychiatric disorders	SCID-5-RV, BDI-II, NEQ

Table 1 - SCID-5-RV – Structured Clinical Interview for the DSM-5 Research Version, EDE-BSV – Eating Disorder Examination – Bariatric Surgery Version, BES - Binge-Eating Depression Scale, HADS - Hospital Depression Scale, BDI-II – Beck Depression Inventory-II, BES – Binge Eating Scale, EDDS – Eating Disorder Diagnostic Scale, EDI – Eating Disorder Inventory, NEQ – Night Eating Questionnaire

4. DISCUSSION

Eating pathology before and after surgery

Binge eating disorder remains the second most common psychiatric disorder in the bariatric population after major depressive disorder (Sarwer et al., 2004). It is characterized by recurrent overeating with the absence of any compensatory weight reduction strategies (e.g., excessive exercises or self-induced vomiting). The sense of lack of control over eating during the episode is also mandatory. Previous research shows that BED prevalence was highest before the surgery (12,7%), with the most significant decrease in the first year (Smith et al., 2019). BED examination post-surgery continues to pose a difficulty as most patients are unable to eat “an amount of food that is definitely larger than most people would eat in a similar period”, as stated in DSM-V, due to the limited capacity of the gastric pouch or sleeve, (Meany et al., 2014). The newest research emphasizes this diagnostic flaw by suggesting that researchers should adjust the criteria, thus objectifying “a large portion of food” for patients to meet the diagnostic requirements.

As such, within bariatric surgery literature, this phenomenon is frequently referred to as subjective binge eating (SBE) Conceição et al., (2015), Goldschmidt et al., (2016) and might be the explanation for why studies show the sudden decline in the prevalence of BED in the first months during post-surgery observation, only for this values to then rise in the following years (Smith et al., 2019). The concept of ‘loss of control eating’ (LOCE) emerged as a primary focus of interest, with a prevalence of 35% before surgery (Smith et al., 2019). The term LOCE is now being used at times instead of BED because of the latter limitations. There have also been suggestions that it is a better indicator of psychopathology – the severity of the loss of control episode corresponds with psychological impairment (Ivezaj et

al., 2018; Goldschmidt, 2017). Lydecker et al., (2019) conducted a study exclusively on patients affected by LOCE Furtado et al., (2023) that showed a co-occurrence with other eating disorders – secretive eating.

This phenomenon became a common form of post-surgery behavior, given the stigma of bariatric surgery combined with strict dietary regulations (Phelan, 2018). Additionally, among this group, body dissatisfaction was higher, even compared to those meeting the criteria for BED. However, it was less related to weight regain than those who do not report secretive eating. Both BED and LOCE remain monitored (Table 1) by researchers to identify risk factors for poor weight loss and weight regain after surgery. Previous reviews stated that, essentially, eating disorders *before* surgery are inconclusive prognostics of weight loss and adherence to post-surgery recommendations (Williams-Kerver et al., 2019; Taba et al., 2021). In contrast, there is an increasing number of studies backing the observation that eating pathology *after* the surgery correlates positively with weight regain (Sarwer et al., 2004).

Another concerning topic is the occurrence of other unclassified eating pathologies – grazing, picking, nibbling, eating in the absence of hunger or beyond satiety, or even excessive dietary restrictions (Ivezaj et al., 2021; Conceição and Goldschmidt, 2019). Bariatric surgery literature suggests that currently used questionnaires such as Eating Disorder Examination – Bariatric Surgery Version (EDE-BSV) or Eating Disorder Diagnostic Scale (EDDS) may not be efficient diagnostic tools to diagnose and distinguish between them properly. Additional research is required in this field to understand the importance of formerly mentioned phenomena better; currently used methods lack consensus, and utilizing consistent definitions and assessment methodologies in future studies should be prioritized.

5. CONCLUSION

Collectively, the latest research remains consistent with previous reviews. Among the diverse group of individuals, with different approaches conducted by each study, problematic eating behaviors were associated with poorer adherence to weight loss and quicker weight regain. Undeniable effects of both BED and LOCE in this population, especially post-surgery, can significantly affect the results of bariatric surgery. Future studies may add to bariatric surgery literature by using consistent terminology and definitions when referring to various eating pathologies. Nonetheless, the factors contributing to successful outcomes are complex, and additional research is needed to improve our understanding of emerging complications. For example, long-term medical observation consisting of self-monitoring of weight and food intake combined with professional assessment would help screen patients as their condition worsens.

Acknowledgment

We thank the participants who were all contributed samples to the study.

Author's contribution

Jan Kościan: Project administration, Supervision

Aleksander Górny: Conceptualization, Investigation

Justyna Chwiejczak: Methodology, Investigation

Michał Obrębski: Data Curation, Formal analysis

Anna Seroka: Resources, Visualization

Karolina Szczerkowska: Resources, Visualization

Maria Mitkowska: Writing – Original Draft

Anna Wójcik: Writing – Review & Editing

Julita Młynarska: Formal analysis

Maria Rybicka: Writing – Review & Editing

Zuzanna Dryżałowska: Writing – Review & Editing

Ethical approval

Not applicable.

Informed consent

Not applicable.

Funding

This study has not received any external funding.

Conflict of interest

The authors declare that there is no conflict of interests.

Data and materials availability

All data sets collected during this study are available upon reasonable request from the corresponding author.

REFERENCES

- Allison KC, Wu J, Spitzer JC, McCuen-Wurst C, Ashare RL, Tewksbury C, LaGrotte CA, Wadden TA, Williams NN, Sarwer DB. Changes in Eating Behaviors and Their Relation to Weight Change 6 and 12 Months After Bariatric Surgery. *Obes Surg* 2023; 33(3):733-742. doi: 10.1007/s11695-022-06442-w
- Athanasiadis DI, Martin A, Kapsampelis P, Monfared S, Stefanidis D. Factors associated with weight regain post-bariatric surgery: a systematic review. *Surg Endosc* 2021; 35(8):4069-4084. doi: 10.1007/s00464-021-08329-w
- Brown W, Anvar M, Liem R, Boza C, Al-Sabah S. 8th IFSO Global Registry Report. IFSO 2023.
- Conceição EM, Goldschmidt A. Disordered eating after bariatric surgery: clinical aspects, impact on outcomes, and intervention strategies. *Curr Opin Psychiatry* 2019; 32(6):504-509. doi: 10.1097/YCO.0000000000000549
- Conceição EM, Utzinger LM, Pisetsky EM. Eating Disorders and Problematic Eating Behaviours Before and After Bariatric Surgery: Characterization, Assessment and Association with Treatment Outcomes. *Eur Eat Disord Rev* 2015; 23(6):417-25. doi: 10.1002/erv.2397
- Courcoulas AP, King WC, Belle SH, Berk P, Flum DR, Garcia L, Gourash W, Horlick M, Mitchell JE, Pomp A, Pories QJ, Purnell JQ, Singh A, Spaniolas K, Thirbly R, Wolfe BM, Yanovski SZ. Seven-Year Weight Trajectories and Health Outcomes in the Longitudinal Assessment of Bariatric Surgery (LABS) Study. *JAMA Surg* 2018; 153(5):427-34. doi: 10.1001/jamasurg.2017.5025
- Furtado TA, Girundi MG, Campolina COC, Mafra SC, Oliveira AMO, Santos MLPDD, Lopes SF, Freire MA. Depressive and eating disorders in patients post-bariatric surgery with weight regain: a descriptive observational study. *Arq Bras Cir Dig* 2023; 36:e1725. doi: 10.1590/0102-672020230002e1725
- Goldschmidt AB, Conceição EM, Thomas JG, Mitchell JE, Raynor HA, Bond DS. Conceptualizing and studying binge and loss of control eating in bariatric surgery patients—time for a paradigm shift? *Surg Obes Relat Dis* 2016; 12(8):1622-1625. doi: 10.1016/j.soard.2016.09.008
- Goldschmidt AB. Are loss of control while eating and overeating valid constructs? A critical review of the literature. *Obes Rev* 2017; 18(4):412-449. doi: 10.1111/obr.12491
- Hassapidou M, Vlassopoulos A, Kalliostra M, Govers E, Mulrooney H, Ells L, Salas XR, Muscogiuri G, Darleska TH, Busetto L, Yumuk VD, Dicker D, Halford J, Woodward E, Douglas P, Brown J, Brown T. European Association for the Study of Obesity Position Statement on Medical Nutrition Therapy for the Management of Overweight and Obesity in Adults Developed in Collaboration with the European Federation of the Associations of Dietitians. *Obes Facts* 2023; 16(1):11-28. doi: 10.1159/000528083
- Ivezaj V, Barnes RD, Cooper Z, Grilo CM. Loss-of-control eating after bariatric/sleeve gastrectomy surgery: Similar to binge-eating disorder despite differences in quantities. *Gen Hosp Psychiatry* 2018; 54:25-30. doi: 10.1016/j.genhosppsych.2018.07.002
- Ivezaj V, Carr MM, Brode C, Devlin M, Heinberg LJ, Kalarchian MA, Sysko R, Williams-Kerver G, Mitchell JE. Disordered eating following bariatric surgery: a review of measurement and conceptual considerations. *Surg Obes Relat Dis* 2021; 17(8):1510-1520. doi: 10.1016/j.soard.2021.03.008
- Taba JV, Suzuki MO, Nascimento FSD, Iuamoto LR, Hsing WT, Pipek LZ, Carneiro-D'Albuquerque LA, Meyer A, Andraus W. The Development of Feeding and Eating Disorders after Bariatric Surgery: A Systematic Review and Meta-Analysis. *Nutrients* 2021; 13(7):2396. doi: 10.3390/nu13072396

14. Kloock S, Ziegler CG, Dischinger U. Obesity and its comorbidities, current treatment options and future perspectives: Challenging bariatric surgery? *Pharmacol Ther* 2023; 251:108549. doi: 10.1016/j.pharmthera.2023.108549
15. Law S, Dong S, Zhou F, Zheng D, Wang C, Dong Z. Bariatric surgery and mental health outcomes: an umbrella review. *Front Endocrinol (Lausanne)* 2023; 14:1283621. doi: 10.3389/fendo.2023.128362
16. Lobstein T, Jackson-Leach R, Powis J, Brinsden H, Gray M. *World Obesity Atlas 2023*. World Obesity Federation, London, 2023.
17. Lydecker JA, Ivezaj V, Grilo CM. Secretive eating and binge eating following bariatric surgery. *Int J Eat Disord* 2019; 52(8): 935-940. doi: 10.1002/eat.23089
18. Meany G, Conceição E, Mitchell JE. Binge Eating, Binge Eating Disorder and Loss of Control Eating: Effects on Weight Outcomes after Bariatric Surgery. *Eur Eat Disord Rev* 2014; 22(2):87-91. doi: 10.1002/erv.2273
19. Niego SH, Kofman MD, Weiss JJ, Geliebter A. Binge eating in the bariatric surgery population: A review of the literature. *Int J Eat Disord* 2007; 40(4):349-59. doi: 10.1002/eat.20376
20. Noria SF, Shelby RD, Atkins KD, Nguyen NT, Gadde KM. Weight Regain After Bariatric Surgery: Scope of the Problem, Causes, Prevention, and Treatment. *Curr Diab Rep* 2023; 23(3):31-42. doi: 10.1007/s11892-023-01498-z
21. O'Brien PE, Hindle A, Brennan L, Skinner S, Burton P, Smith A, Crosthwaite G, Brown W. Long-Term Outcomes After Bariatric Surgery: a Systematic Review and Meta-analysis of Weight Loss at 10 or More Years for All Bariatric Procedures and a Single-Centre Review of 20-Year Outcomes After Adjustable Gastric Banding. *Obes Surg* 2019; 29(1):3-14. doi: 10.1007/s11695-018-3525-0
22. Obeso-Fernández J, Millan-Alanis JM, Rodríguez-Bautista M, Medrano-Juarez S, Oyervides-Fuentes S, Gonzalez-Cruz D, González-González JG, Rodríguez-Gutiérrez R. Benefits of bariatric surgery on microvascular outcomes in adult patients with type 2 diabetes: a systematic review and meta-analysis. *Surg Obes Relat Dis* 2023; 19(8):916-927. doi: 10.1016/j.soard.2023.02.024
23. Opozda M, Chur-Hansen A, Wittert G. Changes in problematic and disordered eating after gastric bypass, adjustable gastric banding and vertical sleeve gastrectomy: a systematic review of pre-post studies. *Obes Rev* 2016; 17(8):770-92. doi: 10.1111/obr.12425
24. Phelan SM. An update on research examining the implications of stigma for access to and utilization of bariatric surgery. *Curr Opin Endocrinol Diabetes Obes* 2018; 25(5):321-325. doi: 10.1097/MED.0000000000000431
25. Sarwer DB, Cohn NI, Gibbons LM, Magee L, Crerand CE, Raper SE, Rosato EF, Williams NN, Wadden TA. Psychiatric Diagnoses and Psychiatric Treatment Among Bariatric Surgery Candidates. *Obes Surg* 2004; 14(9):1148-56. doi: 10.1381/0960892042386922
26. Sarwer DB, Heinberg LJ. A review of the psychosocial aspects of clinically severe obesity and bariatric surgery. *Am Psychol* 2020; 75(2):252-264. doi: 10.1037/amp0000550
27. Sarwer DB, Wadden TA, Ashare RL, Spitzer JC, McCuen-Wurst C, LaGrotte C, Williams NN, Edwards M, Tewksbury C, Wu J, Wajeu G, Allison KC. Psychopathology, disordered eating, and impulsivity in patients seeking bariatric surgery. *Surg Obes Relat Dis* 2021; 17(3):516-524. doi: 10.1016/j.soard.2020.11.005
28. Smith KE, Orcutt M, Steffen KJ, Crosby RD, Cao L, Garcia L, Mitchell JE. Loss of Control Eating and Binge Eating in the 7 Years Following Bariatric Surgery. *Obes Surg* 2019; 29(6):1773-1780. doi: 10.1007/s11695-019-03791-x
29. White MA, Kalarchian MA, Masheb RM, Marcus MD, Grilo CM. Loss of Control Over Eating Predicts Outcomes in Bariatric Surgery Patients. *J Clin Psychiatry* 2010; 71(2):175-84. doi: 10.4088/JCP.08m04328blu
30. Williams-Kerver GA, Steffen KJ, Mitchell JE. Eating Pathology After Bariatric Surgery: an Updated Review of the Recent Literature. *Curr Psychiatry Rep* 2019; 21(9):86. doi: 10.1007/s11920-019-1071-7