

Medical Science

To Cite:

Józefiak A, Szczepanik M, Bochyński C, Hałasiński P, Kropidłowska D, Horbaczewski M, Piela K, Mazurek J, Mazurek G, Myślicka M, Góralski P, Włodarczyk K. Diagnostic and treatment difficulties in a patient with anemia: Review with Glanzmann's Thrombasthenia. *Medical Science* 2024; 28: e99ms3403

doi: <https://doi.org/10.54905/disssi.v28i150.e99ms3403>

Authors' Affiliation:

¹Department of Internal Diseases and Endocrinology, Gabriel Narutowicz Municipal Specialist Hospital in Cracow, Prądnicka 35-37, 31-202 Cracow, Poland

²Department of Pediatrics, District Hospital in Chrzanów, Topolowa 16, 32-500 Chrzanów, Poland

³Department of Internal Medicine, Railway Hospital. Włodzimierz Roefler, MD, PhD in Pruszków, Warszawa 1, 05-800 Pruszków, Poland

⁴Clinical Department of Gastroenterology, Metabolic, Internal Diseases and Dietetics, University Clinical Hospital in Poznań, 49 Przybyszewskiego, 60-355 Poznań, Poland

⁵Polish Red Cross Maritime Hospital in Gdynia, Powstania Styczniowego 1, 81-519 Gdynia, Poland

⁶Hospital Emergency Department, District Health Center Kartuszy, Floriana Ceynowy 7, 83-300 Kartuszy, Poland

⁷Hospital Emergency Department, University Clinical Hospital. Fryderyk Chopin in Rzeszów, Fryderyka Szopena 2, 35-055 Rzeszów, Poland

⁸Primary Health Care, Private health care facility "ALMUS" Jacek Pieniazek, Białoboki 137, 37-207 Gać, Poland

⁹Department of Cardiology and Internal Diseases, Gabriel Narutowicz Municipal Specialist Hospital in Cracow, Prądnicka 35-37, 31-202 Cracow, Poland

¹⁰Wrocław Medical University, wyb. Ludwika Pasteura 1, 50-367 Wrocław, Poland

¹¹University Clinical Center in Gdańsk, Dębinki 7, 80-952 Gdańsk, Poland

*Corresponding Author

Department of Internal Diseases and Endocrinology, Gabriel Narutowicz Municipal Specialist Hospital in Cracow, Prądnicka 35-37, 31-202 Cracow, Poland

Email: jozefiakanna@onet.pl

Contact List
 Anna Józefiak jozefiakanna@onet.pl
 Magdalena Szczepanik magdalena.szczepanik97@gmail.com
 Cezary Bochyński cezary102@gmail.com
 Przemysław Hałasiński przemek0215@gmail.com
 Dominika Kropidłowska dominikakropidlovska@gmail.com
 Maciej Horbaczewski domenodan@gmail.com
 Kinga Piela piela.kinga@t1@gmail.com
 Jolanta Mazurek M.jasminaj@gmail.com
 Gabriela Mazurek gabriela.mazurek@gmail.com
 Maria Myślicka maria.myślicka36@gmail.com
 Patryk Góralski patryk32150@gmail.com
 Klaudia Włodarczyk wlodarczyk.klaudia@gmail.com

ORCID List
 Anna Józefiak 0000-0003-3968-473X
 Magdalena Szczepanik 0000-0001-5550-0860
 Cezary Bochyński 0009-0001-2672-2445
 Przemysław Hałasiński 0000-0002-7223-3394
 Dominika Kropidłowska 0009-0001-2508-2821
 Maciej Horbaczewski 0009-0002-1842-0591
 Kinga Piela 0009-0002-1507-8065
 Jolanta Mazurek 0009-0001-8484-4387
 Gabriela Mazurek 0000-0001-7338-8628
 Maria Myślicka 0009-0000-6190-9128
 Patryk Góralski 0009-0008-1831-0775
 Klaudia Włodarczyk 0009-0002-0779-8564

Peer-Review History

Received: 12 May 2024

Reviewed & Revised: 16/May/2024 to 02/August/2024

Accepted: 06 August 2024

Published: 16 August 2024

Peer-review Method

External peer-review was done through double-blind method.

Medical Science

pISSN 2321-7359; eISSN 2321-7367



© The Author(s) 2024. Open Access. This article is licensed under a [Creative Commons Attribution License 4.0 \(CC BY 4.0\)](https://creativecommons.org/licenses/by/4.0/), which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>.



Diagnostic and treatment difficulties in a patient with anemia: Review with Glanzmann's Thrombasthenia

Anna Józefiak^{1*}, Magdalena Szczepanik², Cezary Bochyński³, Przemysław Hałasiński⁴, Dominika Kropidłowska⁵, Maciej Horbaczewski⁶, Kinga Piela⁷, Jolanta Mazurek⁸, Gabriela Mazurek⁹, Maria Myślicka¹⁰, Patryk Góralski¹¹, Klaudia Włodarczyk¹⁰

ABSTRACT

Introduction: Glanzmann's thrombasthenia (GT) is a rare disease inherited in an autosomal recessive manner. Qualitative or quantitative defect of the adhesive protein that binds to the platelet receptor in the coagulation process causes this disease. In most cases, symptoms appear in early childhood. **The aim:** This study aims to present the diagnostic and therapeutic problem in a patient with GT. **Case report:** A 70-year-old patient hospitalized due to symptomatic anemia. Initially the patient was diagnosed with hemophilia, later the diagnosis was changed to Glanzmann thrombasthenia. The patient's last hospitalization was in 2024, during which gastroscopy and colonoscopy was performed. No pathological changes were detected. **Results:** Glanzmann thrombasthenia is characterized by an average platelet count and morphology, standard prothrombin time, and standard partial thromboplastin time, accompanied by prolonged bleeding time. The diagnosis is usually based on flow cytometry, which shows the absence of GPIIb or GPIIIa molecules on the platelet surface. **Conclusions:** Glanzmann thrombasthenia (GT) is a rare genetic disease that causes symptoms of bleeding diathesis. Quick diagnosis and appropriate treatment are crucial for managing this condition. Treatment options include transfusions of platelet concentrate, antifibrinolytic drugs, recombinant factor VIIa, and hematopoietic cell transplantation.

Keywords: Glanzmann's thrombasthenia, Bleeding, diagnosis, treatment, inherited thrombocytopathy

1. INTRODUCTION

The ability of platelets to adhere to the subendothelial layer of the vascular epithelium layer and aggregate enables the formation of a hemostatic cap. Under physiological conditions, platelets maintain the integrity of blood vessels (Choińska et al., 2016). Platelet function abnormalities divide into primary (congenital) and secondary (acquired). Primary defects are rare. Acquired platelet abnormalities are far more common and often occur in chronic diseases. There are several classifications of congenital thrombocytopathies. Based on the clinical course, it is possible to distinguish severe forms, which include Glanzmann's thrombasthenia. Glanzmann's thrombasthenia is a very rare thrombotic diathesis with a congenital disability in platelet function (Grassetto et al., 2017).

The synthesis of the Glycoprotein IIb/IIIa complex (GPIIb/IIIa) is impaired, which is a receptor for fibrinogen, fibronectin, Von Willebrand factor, and thrombospondin (Noris and Pecci, 2017; Nurden et al., 2015; Nurden, 2017). GT is a genetic autosomal recessive disease. Patients present with symptoms of haemorrhagic diathesis, even during childhood, e.g. from the gingiva, nose, gastrointestinal tract (Jiménez-Castillo et al., 2023). Prolonged post-traumatic and post-operative bleeding is characteristic (Botero et al., 2020; Ahammad et al., 2020). The treatment of Glanzmann's Thrombasthenia is symptomatic. In patients with severe platelet defects, platelet concentrate is transfused (Nurden, 2019; Poon et al., 2015; Poon et al., 2016; Poon et al., 2023; Balduini, 2022; Rodeghiero, 2022). In our work, we will present the case of a patient suffering from Glanzmann's Thrombasthenia.

2. METHODOLOGY

This case report was conducted by searching for current papers on PubMed and Google Scholar using the search phrases (Glanzmann's thrombasthenia) AND (inherited thrombocytopathy) AND (Bleeding) AND (treatment). After eliminating duplicates, we appraised all publications using the titles and abstracts. Following an exact revision of complete manuscripts, 18 articles met the inclusion criteria. The research took place in April 2024.

3. CASE DESCRIPTION

A 70-year-old patient with Glanzman's thrombasthenia diagnosed about 15 years ago, previously diagnosed with hemophilia, repeatedly hospitalized at the Gabriel Narutowicz City Hospital in Krakow, Department of Internal Medicine and Endocrinology, for transfusions of blood products, the last time in April 2024, currently readmitted for symptomatic anemia with diffuse ecchymoses on legs (Figure 1). The patient negates symptoms of bleeding. On admission, average condition, conscious, in logical contact, weakened, with exertional dyspnea, lasting for two days. There are no features of bleeding on physical examination. Patient with concomitant chronic diseases; life-threatening anemia, hypertension, cholelithiasis without cholecystitis, diaphragmatic hernia, status after removal of a polyp from the hepatic fold of the colon, history of gastric ulcer disease, esophageal hiatal hernia, abnormal glucose tolerance, spinal disc herniation, convergent strabismus, blindness of the left eye, bleeding nodules.

In laboratory tests, life-threatening anemia with Hb=5.7 g/dl, WBC=13.31 thousand/uL, RBC=2.03 million/uL, HCT=18.7%, Creatinine=76 umol/L, Glucose=7.37 mmol/L, Urea=6.0 mmol/L, CRP=2.5 ml/L, Sodium=137 mmol/L, Potassium=4.2 mmol/L, INR=1.20. On gastroscopy, gullet unchanged. Stomach: Polyps in the pre-pyloric part with a diameter of 4-7mm. Pylorus unobstructed. The jejunum and the extra-duodenal part of the duodenum unchanged. Reverse gastroscopy-single soft small polyps below the pylorus. There were no features of upper gastrointestinal bleeding. Histopathological Description: 1. Location: Pre-pyloric part of the stomach (antrum). 2. Changes: Single erosions. During colonoscopy: Examination of the entire large intestine up to the cecum visualized the Bauhin valve and the base of the appendix. Inferior bowel preparation for the examination, with BBPS classification of 1+1+1= 3/9, sectionally retained food debris.

On the fold opposite the Bauhin valve in the cecum, a polyp about 2 mm in diameter was removed with forceps (1s, Kudo II). A cauliflower-shaped polyp about 15-17 mm in diameter on a thin pedicle was removed with a coagulation snare at a distance of about 22-24 cm from the sphincter (1p, Kudo IV). Other than that, no changes were found in the extent of the fragmentary mucosa. Electrocardiogram: Sinus tachycardia 140/min. Intermediate axis. No fresh ischemic changes. The ward transfused nine units of packed red blood cells and four packs of interim platelet units, achieving clinical improvement and increasing blood morphotic parameters. Treatment administered: 9 units PRBC, four packs IPU, Paracetamol, Prestarium, Amlozek, Emanera, Cyclonamide. Laboratory tests at

discharge: WBC= 4.80 thousand/uL, RBC= 3.53 million/uL, HGB= 9.9 g/dL, HCT= 30.5%, Na= 139 mmol/L, Potassium= 4.6 mmol/L (Table 1). Discharged home in general good condition.



Figure 1 Diffuse ecchymoses in a 70-year-old patient with Glanzmann's thrombasthenia.

Table 1 Table showing morphology changes during hospitalization.

Date	2024.06.26	2024.06.25	2024.06.23	2024.06.21	2024.06.19	
Amount of Blood products transfused	-	+2 units of blood platelets	+4 units of blood platelets +4 units of red blood cells	+2 units of blood platelets +2 units of red blood cells	-	-
	-	-	-	-	-	Units
Changes in morphology after transfusions	-	-	-	-	-	-
WBC	4.80	4.20	5.94	7.21	13.31	109/L
RBC	3.53	3.49	3.03	2.81	2.03	1012/L
HGB	9.9	9.7	8.3	7.7	5.7	g/dL
HCT	30.5	29.9	26.6	24.2	18.7	%
PLT	136	83	121	159	190	109/L

*Source photo, taken from a private gallery, made by the authors in the Emergency Room of the Hospital

4. DISCUSSION

Glanzmann's thrombasthenia is a congenital disease first described by Dr. Eduard Glanzmann in 1918, when he reported a patient with prolonged bleeding time and a lack of platelet aggregation. The absence or improper functioning of glycoprotein IIB/IIIa, which plays a crucial role in binding to fibrinogen and participating in the formation of the platelet plug in the process of hemostasis, causes this disease. In the results of blood tests, the level of platelets is usually average or slightly reduced, but their improper functioning prevents the formation of a proper clot (Solh et al., 2015). Understanding the pathogenesis of Glanzmann's thrombasthenia expands our view of the function of platelets in maintaining hemostasis (Lane et al., 2005). This disease usually presents its first symptoms in childhood, when as a result of trauma, there is a significantly prolonged bleeding time, which raises concern and leads to further diagnostics.

When Glanzmann's thrombasthenia is suspected, other more common causes of prolonged bleeding time should first be ruled out. Laboratory tests report an average platelet count and morphology, standard prothrombin time, and normal partial thromboplastin activation time with prolonged bleeding time (Botero et al., 2020). This disease characterizes excessive tendency to bleed, which appears in early childhood. Most often, these are bleeding from the gums, nose, gastrointestinal tract, hematuria, and possible petechiae on the skin. Post-traumatic and postoperative bleeding time is prolonged, and women may experience significantly increased bleeding from the genital tract. In about 3% of patients, intra-articular bleeding occurs, causing damage and, as a consequence, a significant reduction in the quality of life (Pontara et al., 2014).

The diagnosis of the disease bases on the performance of significantly prolonged bleeding time with normal morphology and platelet count values, reduced or no clot retraction, no platelet aggregation under the influence of ADP, collagen, adrenaline, arachidonic acid, with normal aggregation under the influence of ristocetin in flow cytometry, no GPIIb and (or) GPIIIa molecules on the platelet surface (Botero et al., 2020). Currently, there is no effective cure for Glanzmann's thrombasthenia. Transfusions of platelet concentrate and antifibrinolytic drugs, are still the main health treatments methods. Due to the need for frequent transfusions of blood components, it is best to transfuse them following the recipient's HLA, reducing the risk of alloimmunization. In cases where this treatment is insufficient or alloimmunization has occurred, recombinant factor VIIa (Eptacog alfa) can be used. As a last resort, when other therapeutic options have been exhausted, hematopoietic cell transplantation can be performed (Solh et al., 2015).

5. CONCLUSIONS

Glanzmann's thrombasthenia is a rare disease, classified as a congenital thrombocytopathy. The main symptoms of this disease are similar to the symptoms of haemorrhagic diathesis, such as frequent bleeding from the gums and gastrointestinal tract, and petechiae on the skin. Prolonged bleeding after trauma and surgery affect the quality of life of Glanzmann's thrombasthenic patients. The main problem for people struggling with this disease is getting an early correct diagnosis. Although there is no effective cure for Glanzmann's thrombasthenia, symptomatic treatment is applicable. In the absence of such treatment, physicians use recombinant factor VIIa, while in severe Glanzmann's thrombasthenia, haematopoietic cells are transplanted.

Author's Contribution

Anna Józefiak: Conceptualization, methodology, investigation

Magdalena Szczepanik: Conceptualization, methodology, investigation

Cezary Bochyński: Methodology, Review and editing

Przemysław Hałasiński: Conceptualization, writing- rough preparation

Dominika Kropidłowska: Resources, writing- rough preparation

Maciej Horbaczewski: Conceptualization, writing- rough preparation

Kinga Piela: Formal analysis, supervision

Jolanta Mazurek: Review and editing, supervision

Gabriela Mazurek: Visualization, data curation

Maria Myślicka: Formal analysis, supervision

Patryk Góralski: Resources, writing- rough preparation

Klaudia Włodarczyk: Visualization, data curation

Project administration: Cezary Bochyński

Informed consent

Not applicable.

Ethical approval

Not applicable.

Funding

This study has not received any external funding.

Conflict of interest

The authors declare that there is no conflict of interests.

Data and materials availability

All data sets collected during this study are available upon reasonable request from the corresponding author.

REFERENCES

1. Ahammad J, Kamath A, Shastry S, Chitlur M, Kurien A. Clinico-hematological and thromboelastographic profiles in glanzmann's thrombasthenia. *Blood Coagul Fibrinolysis* 2020; 31(1):29-34. doi: 10.1097/MBC.0000000000000870
2. Balduini CL. Treatment of inherited thrombocytopenias. *Haematologica* 2022; 107(6):1278-1292. doi: 10.3324/haematol.2022.280856
3. Botero JP, Lee K, Branchford BR, Bray PF, Freson K, Lambert MP, Luo M, Mohan S, Ross JE, Bergmeier W, Di-Paola J; ClinGen Platelet Disorder Variant Curation Expert Panel. Glanzmann thrombasthenia: genetic basis and clinical correlates. *Haematologica* 2020; 105(4):888-894. doi: 10.3324/haematol.2018.214239

4. Choińska B, Łaguna P, Matysiak M. Trombastenia Glanzmanna– obraz kliniczny, diagnostyka, postępowanie. © Borgis Nowa Pediatria 2016; 20(4):179-182..
5. Grassetto A, Fullin G, Lazzari F, Panizzo F, Polese F, Gessori G, Farnia A. Perioperative ROTEM and ROTEMplatelet monitoring in a case of Glanzmann's thrombasthenia. Blood Coagul Fibrinolysis 2017; 28(1):96-99. doi: 10.1097/MBC.0000000000000532
6. Jiménez-Castillo RA, Benavides-Salgado DE, Jáquez-Quintana JO, González González JA, Cortez Hernández CA, Náñez-Terrerros H, Maldonado-Garza HJ. Glanzmann thrombasthenia: an uncommon cause of acute upper gastrointestinal bleeding. Rev Esp Enferm Dig 2023; 115(3):149. doi: 10.17235/reed.2022.9081/2022
7. Lane DA, Philippou H, Huntington JA. Directing thrombin. Blood 2005; 106(8):2605-12. doi: 10.1182/blood-2005-04-1710
8. Noris P, Pecci A. Hereditary thrombocytopenias: a growing list of disorders. Hematology Am Soc Hematol Educ Program 2017; 2017(1):385-399. doi: 10.1182/asheducation-2017.1.385
9. Nurden AT, Pillois X, Fiore M, Alessi MC, Bonduel M, Dreyfus M, Goudemand J, Gruel Y, Benabdallah-Guerida S, Latger-Cannard V, Négrier C, Nugent D, Oiron RD, Rand ML, Sié P, Trossaert M, Alberio L, Martins N, Sirvain-Trukniewicz P, Couloux A, Canault M, Fronthoth JP, Fretigny M, Nurden P, Heilig R, Vinciguerra C. Expanding the Mutation Spectrum Affecting α IIb β 3 Integrin in Glanzmann Thrombasthenia: Screening of the ITGA2B and ITGB3 Genes in a Large International Cohort. Hum Mutat 2015; 36(5):548-61. doi: 10.1002/humu.22776
10. Nurden AT. Acquired Glanzmann thrombasthenia: From antibodies to anti-platelet drugs. Blood Rev 2019; 36:10-22. doi: 10.1016/j.blre.2019.03.004
11. Nurden AT. Should studies on Glanzmann thrombasthenia not be telling us more about cardiovascular disease and other major illnesses? Blood Rev 2017; 31(5):287-299. doi: 10.1016/j.blre.2017.03.005
12. Pontara E, Gresele P, Cattini MG, Daidone V, Barbon G, Girolami A, Zanon E, Casonato A. Spontaneous hemarthrosis in combined Glanzmann thrombasthenia and type 2N von Willebrand disease. Blood Coagul Fibrinolysis 2014; 25(4):401-4. doi: 10.1097/MBC.0000000000000067
13. Poon MC, Di-Minno G, d'Oiron R, Zotz R. New Insights into the Treatment of Glanzmann Thrombasthenia. Transfus Med Rev 2016; 30(2):92-9. doi: 10.1016/j.tmr.2016.01.001
14. Poon MC, d'Oiron R, Zotz RB, Bindslev N, Di-Minno MN, Di-Minno G. The Glanzmann Thrombasthenia Registry: safety of platelet therapy in patients with Glanzmann thrombasthenia and changes in alloimmunization status. Haematologica 2023; 108(10):2855-2858. doi: 10.3324/haematol.2022.281973
15. Poon MC, d'Oiron R, Zotz RB, Bindslev N, Di-Minno MN, Di-Minno G; Glanzmann Thrombasthenia Registry Investigators. The international, prospective Glanzmann Thrombasthenia Registry: treatment and outcomes in surgical intervention. Haematologica 2015; 100(8):1038-44. doi: 10.3324/haematol.2014.121384
16. Rodeghiero F. Introduction to a review series on the treatment of thrombocytopenic disorders: something old, something new. Haematologica 2022; 107(6):1239-1242. doi: 10.3324/haematol.2022.280920
17. Solh T, Botsford A, Solh M. Glanzmann's thrombasthenia: pathogenesis, diagnosis, and current and emerging treatment options. J Blood Med 2015; 8:219-227. doi: 10.2147/JBM.S71319